

On the local zeta functions and the b -functions of certain hyperplane arrangements

(with Appendix by Willem Veys)

Nero Budur, Morihiko Saito, and Sergey Yuzvinsky

Abstract. Conjectures of J. Igusa for p -adic local zeta functions and of J. Denef and F. Loeser for topological local zeta functions assert that (the real part of) the poles of these local zeta functions are roots of the Bernstein-Sato polynomials (i.e. the b -functions). We prove these conjectures for certain hyperplane arrangements, including the case of reduced hyperplane arrangements in three-dimensional affine space.

Introduction

Let K be a p -adic field, i.e. a finite extension of $\mathbf{Q}_p$, and $\mathcal{O}_K$ be the ring of integers of K . We have the norm defined by $|x|_K = q^{-v(x)}$ for $x \in K^*$ where $v(x) \in \mathbf{Z}$ is the valuation (or the order) of $x \in K$ and q is the cardinality of the residue field $\mathcal{O}_K/\mathfrak{m}_K$ with $\mathfrak{m}_K$ the maximal ideal. For a nonconstant polynomial $f \in K[x_1, \dots, x_n]$, Igusa's p -adic local zeta function (associated with the characteristic function of $\mathcal{O}_K^n \subset K^n$, see [Ig1], [Ig4]) is defined by the meromorphic continuation of the integral

$$Z_f^p(s) = \int_{\mathcal{O}_K^n} |f(x)|_K^s dx \quad \text{for } \operatorname{Re} s > 0.$$

Here dx denotes the Haar measure on the compact open subgroup $\mathcal{O}_K^n$ of K^n , which is the p -adic analogue of the polydisk Δ^n in $\mathbf{C}^n$. Note that $Z_f^p(s)$ is closely related to the Poincaré series associated with the numbers of solutions of $f = 0$ in $(\mathcal{O}_K/\mathfrak{m}_K^i)^n$ for $i > 0$ in the case $f \in \mathcal{O}_K[x_1, \dots, x_n]$.

On the other hand, the Bernstein-Sato polynomial (i.e. the b -function) of a polynomial $f \in K[x_1, \dots, x_n]$ is the monic polynomial $b_f(s)$ of the least degree satisfying the relation

$$b_f(s)f^s = Pf^{s+1} \quad \text{in } R_f[s]f^s \quad \text{for some } P \in \mathcal{D}_n[s],$$

where R_f is the localization of $R := K[x_1, \dots, x_n]$ by f and $\mathcal{D}_n$ is the Weyl algebra which is generated over K by $x_1, \dots, x_n$ and $\partial/\partial x_1, \dots, \partial/\partial x_n$. Here K may be any field of

2000 Mathematics Subject Classification: 14J17, 32S40, 32S22.

The first author is supported by the NSF grant DMS-0700360.

The second author is partially supported by Kakenhi 21540037.

characteristic 0, and $b_f(s)$ is invariant by extensions of the field K , see (2.1) below. (There is a shift of the variable s by 1 if one uses the definition of the Bernstein polynomial in [Be] since f^s is replaced by f^{s-1} there). The local b -function $b_{f,x}(s)$ is defined by replacing the Weyl algebra $\mathcal{D}_n$ with $\mathcal{D}_{X,x}$. Note that for a homogeneous polynomial f , we have $b_f(s) = b_{f,0}(s)$.

A conjecture of J. Igusa [Ig2] asserts the following.

Conjecture (A)^p. The real part of any pole of the p -adic local zeta function $Z_f^p(s)$ is a root of $b_f(s)$.

Inspired by this conjecture, J. Denef and F. Loeser [DL] defined the topological local zeta function $Z_{f,x}^{top}(s)$ (see (1.1.1) below) for a nonconstant polynomial f and $x \in f^{-1}(0)$ in the case $K = \mathbf{C}$, and conjectured the following.

Conjecture (A)^{top}. Any pole of the topological local zeta function $Z_{f,x}^{top}(s)$ is a root of $b_{f,x}(s)$.

There is a weaker version of the conjectures, due to Igusa, and Denef and Loeser respectively, and called the *monodromy conjecture*, as follows.

Conjecture (B)^p. For any pole α of the p -adic local zeta function $Z_f(s)$, $e^{2\pi i \operatorname{Re}(\alpha)}$ is an eigenvalue of the Milnor monodromy of $f_{\mathbf{C}}$ at some $x \in f_{\mathbf{C}}^{-1}(0) \subset \mathbf{C}^n$ choosing an embedding $K \hookrightarrow \mathbf{C}$, where $f_{\mathbf{C}}$ is the image of f in $\mathbf{C}[x_1, \dots, x_n]$.

Conjecture (B)^{top}. For any pole α of the topological local zeta function $Z_{f,x}^{top}(s)$, $e^{2\pi i \alpha}$ is an eigenvalue of the Milnor monodromy of f at $y \in f^{-1}(0)$ sufficiently near x .

In Conjecture (B)^p, it is enough to consider an embedding $K^f \hookrightarrow \mathbf{C}$, where K^f is the subfield of K generated by the coefficients of the linear factors of f so that D is defined over K_f . Originally Conjecture (A)^p and (B)^p are stated for a polynomial $f \in F[x_1, \dots, x_n]$ with F a number field and K the completion of F at a prime of F (except possibly for a finite number of primes). In the hyperplane arrangement case, however, this assumption does not seem to be essential since Conjecture (B)^p is already proved by [BMT] and Conjecture (A)^p is reduced to Conjecture (C) below.

By Conjecture (A) we will mean Conjecture (A)^p or Conjecture (A)^{top} depending on whether K is the p -adic or complex number field, and similarly for Conjecture (B). Note that the eigenvalues of the Milnor monodromies in Conjecture (B) can be defined in a purely algebraic way using the V -filtration of Kashiwara [Ka2] and Malgrange [Ma2] on the $\mathcal{D}_n[s]$ -module $R_f[s]f^s$, and the union of the eigenvalues of the Milnor monodromies for $x \in f_{\mathbf{C}}^{-1}(0) \subset \mathbf{C}^n$ is independent of the choice of the embedding $K^f \hookrightarrow \mathbf{C}$, see (2.1) below. Moreover we have the following.

Proposition 1. *Let K be a subfield of $\mathbf{C}$, and $f \in K[x_1, \dots, x_n]$.*

(i) *A complex number $\lambda \in \mathbf{C}$ is an eigenvalue of the Milnor monodromy of $f_{\mathbf{C}}$ at some $x \in f_{\mathbf{C}}^{-1}(0) \subset \mathbf{C}^n$ if and only if there is a root α of $b_f(s)$ such that $\lambda = e^{-2\pi i \alpha}$.*

(ii) *If $K = \mathbf{C}$, then for any $x \in f^{-1}(0)$, there is an open neighborhood U of x in classical topology such that for any open neighborhood U' of x in U , the following two conditions are equivalent.*

- (a) The number λ is an eigenvalue of the Milnor monodromy of f at some $y \in f^{-1}(0) \cap U'$.
- (b) There is a root α of $b_{f,x}(s)$ such that $\lambda = e^{-2\pi i \alpha}$.

This follows from [Ka1], [Ma2]. By Proposition 1, Conjecture (B) can be viewed as the modulo $\mathbf{Z}$ version of Conjecture (A), and is weaker than the latter. It is known that Conjectures (A) and (B) are rather difficult to prove, see e.g. [ACLM1], [ACLM2], [Den], [DL], [Ig3], [Ig4], [KSZ], [Lo1], [Lo2], [LVa], [LV1], [LV2], [Ro], [VP], [Ve1], [Ve2], [Ve3], [Ve4]. For a generalization to the ideal case, see [HMY], [VV] (using [BMS]).

In this paper we prove Conjecture (A) for certain affine hyperplane arrangements D in K^n . Let D_i be the irreducible components of D , and m_i be the multiplicity of D along D_i . Let f be a defining equation of D . Set $d := \deg D = \deg f = \sum_i m_i$. In [BMT], Conjecture (A) is reduced to the following.

Conjecture (C). Let D be an indecomposable essential central hyperplane arrangement in $\mathbf{C}^n$ with degree d . Then $b_f(-n/d) = 0$.

Here central and essential respectively mean that $0 \in D_i$ for any i and $\dim \bigcap_i D_i = 0$. We say that D is *indecomposable* if it is not a union of the pullbacks of arrangements by the two projections of some decomposition $\mathbf{C}^n = \mathbf{C}^{n'} \times \mathbf{C}^{n''}$ as a vector space. Note that the proof of Conjecture (B) in [BMT] implies that $-n/d - 1$ is a root of $b_f(s)$ in case $-n/d$ is not, since the roots of $b_f(s)$ are in $(-2, 0)$, see [Sa1].

As for the reduction of Conjecture (A) to Conjecture (C) we have more precisely the following.

Theorem 1 [BMT]. *For an affine hyperplane arrangement D in K^n , Conjecture (A) holds if Conjecture (C) for $(D/L)_{\mathbf{C}}$ holds for every dense edge L of D .*

Here an edge means an intersection of D_i , and D/L denotes the arrangement in K^n/L defined by the D_i containing L and with the same multiplicity m_i , where we may assume $0 \in L$ replacing the origin of K^n if necessary. We call an edge $L \neq K^n$ *dense* if D/L is indecomposable. If K is a p -adic field, then $(D/L)_{\mathbf{C}}$ denotes the scalar extension of D/L defined by choosing an embedding $K^f \hookrightarrow \mathbf{C}$ where $K^f \subset K$ is the smallest subfield such that f and all the D_i are defined over K^f . We have $(D/L)_{\mathbf{C}} = D/L$ in the case $K = \mathbf{C}$.

Theorem 1 is proved by using a resolution of singularities obtained by blowing up only the proper transforms of the *dense* edges in [STV] (together with Igusa's calculation of candidates for poles of the p -adic zeta functions [Ig1] in the p -adic case, see also (1.1.3) below). Because of this very special kind of resolution, all the obtained candidates for poles contribute at least to the monodromy eigenvalues, and Conjecture (B) is proved in [BMT] for all the candidates for poles using the calculation of the Milnor cohomology of hyperplane arrangements in [CS] (or [Di], Prop. 6.4.6) together with a result of [STV] on the relation between indecomposability and nonvanishing of the Euler characteristic of the projective complement. This is contrary to the most other cases where lots of cancelations of apparent poles occur, see [Den], [Lo1], [Ve1], [Ve2], [Ve3] (and Remark (1.2) below). Recently W. Veys informed us that there are examples of hyperplane arrangement of degree d in $\mathbf{C}^n$ such that $-n/d$ is not a pole of $Z_{f,0}^{top}(s)$ in the case $n = 3$ with D non-reduced or $n = 5$ with D reduced, see Appendix. These examples imply a negative answer to Question (Q) in (1.4). There are no such examples if $n = 2$ or $n = 3$ with D reduced, see Propositions (1.5) and (1.8) below.

In this paper we prove the following.

Theorem 2. *Conjecture (C) holds in the following cases.*

- (i) $\{0\}$ is a good dense edge of D .
- (ii) D is reduced with $n \leq 3$.
- (iii) D is reduced, $(n, d) = 1$, and D_d is generic relative to the other D_j ($j \neq d$).

Here L is called a *good dense edge* if for any dense edges $L' \supset L$, we have

$$n(L)/d(L) \leq n(L')/d(L'),$$

where $d(L) = \text{mult}_L D = \sum_{D_i \supset L} m_i$ and $n(L) = \text{codim } L$. We say that D_d is *generic relative to the other D_j* ($j \neq d$) if any nonzero intersection of D_j ($j \neq d$) is not contained in D_d , see [FT], Example 4.5.

In case (i), Theorem 2 follows from Teitler's refinement [Te] of Mustață's formula [Mu] for multiplier ideals using only dense edges, together with a well-known relation between the jumping coefficients and the roots of $b_f(s)$, see [ELSV]. In case (ii) or (iii), we use a generalization of Malgrange's formula for the roots of $b_f(s)$ in the isolated singularity case (see [Sa1], [Sa2]) reducing the assertion to a certain combinatorial problem which can be solved under condition (ii) or (iii), where we need a result from [FT] in case (iii).

Combining Theorems 1 and 2, we get

Theorem 3. *For an affine hyperplane arrangement D in K^n , conjecture (A) holds if for every dense edge L of D , one of the three conditions in Theorem 2 is satisfied for $(D/L)_{\mathbf{C}}$. In particular, Conjecture (A) holds in the following cases.*

- (i) D is of moderate type.
- (ii) D is reduced with $n \leq 3$.
- (iii) D is reduced with $n = 4$, and for each 0-dimensional dense edge L of D , either condition (ii) or (iii) in Theorem 2 is satisfied for $(D/L)_{\mathbf{C}}$.

Here D is called *of moderate type* if all the dense edges are good. Note that in the case (iii), condition (ii) in Theorem 2 is satisfied for $(D/L)_{\mathbf{C}}$ with $L \neq 0$. It seems quite difficult to generalize the arguments in this paper to the non-reduced case even for $n = 3$, or to the 4-dimensional case even for reduced D .

We would like to thank W. Veys for useful comments and especially for examples in Appendix solving Question (Q) in (1.4) negatively.

In Section 1 we recall some facts from the theory of local zeta functions. In Section 2 we explain how to calculate the b -functions of homogeneous polynomials, and prove Theorem 2 in cases (i) and (iii). In Section 2 we prove Theorem 2 in case (ii). In Appendix by W. Veys, we describe some examples related to Question (Q) in (1.4).

1. Local zeta functions

1.1. Let K be the complex or p -adic number field. Let X be a complex manifold of dimension n with f a holomorphic function on X if $K = \mathbf{C}$, and $X = K^n$ with $f \in K[x_1, \dots, x_n]$ if K is a p -adic field. Set $D = f^{-1}(0)$. Let $\sigma : (\tilde{X}, E) \rightarrow (X, D)$ be an embedded resolution with E_j the irreducible components of $E := \sigma^*D$. Set

$$E_I^\circ = \bigcap_{i \in I} E_i \setminus \bigcup_{i \notin I} E_i, \quad m_j = \text{mult}_{E_j} \sigma^*D, \quad r_j = \text{mult}_{E_j} \det(\text{Jac}(\sigma)).$$

If $K = \mathbf{C}$, the topological local zeta function for $x \in D$ is defined by

$$(1.1.1) \quad Z_{f,x}^{\text{top}}(s) = \sum_I \chi(E_I^\circ \cap \sigma^{-1}(x)) \prod_{j \in I} \frac{1}{m_j s + r_j + 1},$$

which is independent of the choice of the resolution (see [DL]). So we get candidates for poles

$$(1.1.2) \quad \alpha_j := -\frac{r_j + 1}{m_j}.$$

Note that each α_j is not necessarily a pole of $Z_{f,x}^{\text{top}}(s)$ in general. It is not easy to determine exactly false poles since there are cancelations of poles in many cases, see [Den], [Lo1], [Ve1], [Ve2], [Ve3] (and Remark (1.2) below). In the hyperplane arrangement case, however, there is a special kind of resolution by [STV] so that Conjecture (B) is proved for the above candidates for poles although it is still unclear whether they are really poles.

The situation is similar in the p -adic case where Igusa's calculation (see e.g. [Ig4], Theorem 8.2.1 or [Den]) implies that the poles of the local zeta function are among the complex numbers

$$(1.1.3) \quad \alpha_{j,k} := -\frac{r_j + 1}{m_j} - \frac{2\pi\sqrt{-1}k}{m_j \log q} \quad (k \in \mathbf{Z}).$$

1.2. Remark. It is known that there are remarkable cancelations of poles by the summation in the definition (1.1.1). So it is not easy to eliminate false poles, although the curve case is rather well understood, see [Den], [Lo1], [Ve1], [Ve2], [Ve3]. (For a relatively simple proof of Conjecture (B) for $n = 2$, see [Ro].) It is also known that only a few of the roots of $b_f(s)$ can be detected by the local zeta function.

1.3. Proposition. *Let D be a hyperplane arrangement defined by a polynomial f . Then the topological local zeta function $Z_{f,x}^{\text{top}}(s)$ is a combinatorial invariant.*

Proof. By the definition of $Z_{f,x}^{\text{top}}(s)$ in (1.1.1), we may assume D is central, $x = 0$. We have to calculate the Euler characteristic of each open stratum of a stratification of $\sigma^{-1}(0)$ which is induced from the canonical stratification of a divisor with normal crossings. In this case σ is obtained by taking first the blow-up $X' \rightarrow X = \mathbf{C}^n$ along the origin of $\mathbf{C}^n$,

and then taking the base change of an embedded resolution of $(\mathbf{P}^{n-1}, Z)$ by the projection $X' \rightarrow \mathbf{P}^{n-1}$ where $Z := \mathbf{P}(D)$. The Euler characteristic of an open stratum is calculated from those of the closed strata contained in the closure of the given stratum. So the assertion follows by induction on n using [DP] together with the embedded resolution of $(\mathbf{P}^{n-1}, Z)$ obtained by blowing up along the proper transforms of *all* the edges of Z . Indeed, any intersection of the proper transforms of exceptional divisors can be written as a product of embedded resolutions for certain induced arrangements, see loc. cit. and [BS], Prop. 2.7 in this case. (If we blow up along only the proper transforms of dense edges, we can not apply an inductive arguments since there is a problem as below: For two dense edges $L \subset L'$ of $Z \subset \mathbf{P}^{n-1}$, L is not necessarily a dense edge of the induced arrangement in L' .) This finishes the proof of Proposition (1.3).

1.4. Analogue of Conjecture (C). The following question arises naturally:

Question (Q). Let D be an indecomposable essential central hyperplane arrangement in $\mathbf{C}^n$ defined by a polynomial f of degree d . Then, is $-n/d$ a pole of $Z_{f,0}^{top}(s)$?

We have a positive answer to this question if $n = 2$ or $n = 3$ and D is reduced, see Propositions (1.5) and (1.8) below. Recently, W. Veys informed us that the answer is negative in general, more precisely, if $n = 3$ with D non-reduced or $n = 5$ with D reduced, see Appendix.

Assume, for example, $n = 2$ and $d = \sum_{i=1}^e m_i$ with $m_i = \text{mult}_{D_i} D$. Then

$$(1.4.1) \quad Z_{f,0}^{top}(s) = \frac{1}{ds+2} \left(2 - e + \sum_{i=1}^e \frac{1}{m_i s + 1} \right).$$

This immediately follows from the definition of the zeta function since the embedded resolution is obtained by one blow-up and $2 - e$ is the Euler characteristic of the open stratum in $\mathbf{P}^1$. So $-2/d$ is a pole of order 2 if and only if $2m_i = d$ for some i . If $-2/d$ is not a pole of order 2, then the coefficient $C_{-2/d}$ of $\frac{1}{ds+2}$ is given by

$$C_{-2/d} = 2 - e + \sum_{i=1}^e \frac{d}{d - 2m_i}.$$

The next Proposition gives a positive answer to Question (Q) in (1.4) for $n = 2$ where D may be non-reduced. This is a special case of [Ve3], Prop. 2.8.

1.5. Proposition (W. Veys [Ve3]). *With the above notation, assume $n = 2$. Then $-2/d$ is a pole of $Z_{f,0}^{top}(s)$. More precisely, if $-2/d$ is not a pole of order 2, then $C_{-2/d} > 0$ if $\{0\}$ is a good dense edge of D , and $C_{-2/d} < 0$ otherwise.*

Proof. See [Ve3], Prop. 2.8.

1.6. Proposition. *Assume $n = 3$, and D is reduced. Let ν_m ($m \geq 2$) be the number of points of $Z := \mathbf{P}(D)$ with multiplicity m . Then*

$$Z_{f,0}^{top}(s) = \frac{1}{ds+3} \left(\chi(\mathbf{P}^2 \setminus Z) + \frac{\chi(Z \setminus Z^{\text{sing}})}{s+1} + \sum_m \left(2 - m + \frac{m}{s+1} \right) \frac{\nu_m}{ms+2} \right).$$

In particular, $-3/d$ is the only candidate for the pole of order 2 of $Z_{f,0}^{top}(s)$, and is really a pole of order 2 if and only if $d/3 \in \mathbf{Z}$ and $\nu_{2d/3} \neq 0$. If $-3/d$ is not a pole of order 2, the coefficient $C_{-3/d}$ of $\frac{1}{ds+3}$ is given by

$$C_{-3/d} = \frac{9}{d-3} \left(d-1 + \sum_{m \neq 2d/3} \frac{m(m-1)}{2d-3m} \nu_m \right).$$

Proof. Since the embedded resolution of $(\mathbf{P}^2, Z)$ is obtained by blowing up along the singular points of Z , the first assertion follows from the definition of $Z_{f,0}^{top}(s)$ using the partition of the summation over $m \neq 2$ and $m = 2$. This implies the second assertion since the coefficient of the double pole is given up to a nonzero multiplicative constant by

$$2 - m + \frac{md}{d-3} = \frac{2a-1}{a-1} \neq 0, \text{ where } m = 2a \text{ with } a := d/3 \in \mathbf{Z}.$$

For the simple pole case, we have

$$C_{-3/d} = \chi(\mathbf{P}^2 \setminus Z) + \frac{\chi(Z \setminus Z^{\text{sing}})d}{d-3} + \sum_{m \neq 2d/3} \left(2 - m + \frac{md}{d-3} \right) \frac{\nu_m d}{2d-3m}.$$

Here

$$\begin{aligned} \chi(\mathbf{P}^2 \setminus Z) &= 3 - 2d + \sum_m (m-1)\nu_m, \\ \chi(Z \setminus Z^{\text{sing}}) &= 2d - \sum_m m\nu_m. \end{aligned}$$

Indeed, the first equality is reduced to the calculation of $\chi(Z)$ which is obtained by using the short exact sequence $0 \rightarrow \mathbf{Q}_Z \xrightarrow{\iota} \bigoplus_i \mathbf{Q}_{Z_i} \rightarrow \text{Coker } \iota \rightarrow 0$, since the cokernel of ι is supported on the singular points of Z and its rank at p is $m_p - 1$ where m_p is the multiplicity of Z at p .

Substituting these, we see that $C_{-3/d}$ is given by

$$3 - 2d + \frac{2d^2}{d-3} + \sum_{m \neq 2d/3} \nu_m \left(m-1 - \frac{md}{d-3} + \frac{(2-m)d}{2d-3m} + \frac{md^2}{(d-3)(2d-3m)} \right).$$

After some calculation this is transformed to

$$\frac{9}{d-3} \left(d-1 + \sum_{m \neq 2d/3} \frac{m(m-1)}{2d-3m} \nu_m \right).$$

(The detail is left to the reader.) This finishes the proof of Proposition (1.6).

1.7. Remark. A strong form of the conjecture in [DL] predicts that the multiplicity of each root of the zeta function is at most that of the b -function. In general, the multiplicity of the root -1 of the b -function of a reduced essential central hyperplane arrangement is n (see [Sa2], Th. 1), and this settles the problem for the root -1 . However, the problem

is rather difficult for the roots with multiplicity 2 even in the case $n = 3$. In this case the only such root is $-3/d$ with $d/3 \in \mathbf{N}$ and $\nu_{2d/3} \neq 0$ by Proposition (1.6), but it is not easy to calculate the b -function. (Indeed, the multiplicity is calculated only in the case $\nu_m = 0$ for $m > 3$ in loc. cit.)

Using Proposition (1.6) we get the Proposition below which gives a positive answer to Question (Q) in (1.4) if D is reduced and $n = 3$. W. Veys has informed us that he had verified an analogue of it for the (finer) motivic or Hodge zeta functions. (Here ‘finer’ means that the non-vanishing of the pole for these do not imply that for $Z_{f,0}^{top}(s)$ although the converse is true.)

1.8. Proposition. *Let D be an indecomposable essential central hyperplane arrangement of degree d in $\mathbf{C}^3$. Assume D is reduced. Then $-3/d$ is a pole of $Z_{f,0}^{top}(s)$. More precisely, if $-3/d$ is not a pole of order 2, then the coefficient $C_{-3/d}$ of $\frac{1}{ds+3}$ is strictly positive if $\{0\}$ is a good dense edge of D , i.e. if $m < 2d/3$ for any m with $\nu_m \neq 0$, and $C_{-3/d}$ is strictly negative otherwise.*

Proof. We may assume $n = 3$ since the case $n = 2$ is trivial. We may further assume $\{0\}$ is not a good dense edge of D , since the assertion in the good dense edge case easily follows from Proposition (1.6). We may thus assume $\nu_{m_0} \neq 0$ for some $m_0 := 2a + e$ with $0 < e < a := d/3$ where we do not assume $a \in \mathbf{Z}$. Since the sum of the multiplicities of any two singular points of Z is at most $d + 1$, we have $\nu_{m_0} = 1$ and

$$m \leq 3a - m_0 + 1 = a - e + 1 \quad \text{for any } m \neq m_0 \text{ with } \nu_m \neq 0.$$

By Proposition (1.6) the assertion $C_{-3/d} < 0$ is equivalent to

$$\frac{(2a+e)(2a+e-1)}{e} > 3(3a-1) + \sum_{m \leq a-e+1} \frac{m(m-1)}{2a-m} \nu_m.$$

To show the last inequality, we may replace $\frac{m(m-1)}{2a-m}$ with $\frac{m(m-1)}{a+e-1}$, since $\frac{1}{a+e-1} \geq \frac{1}{2a-m}$ for $m \leq a - e + 1$. Using $\binom{d}{2} = \sum_m \binom{m}{2} \nu_m$, the assertion is then reduced to

$$\frac{(2a+e)(2a+e-1)}{e} > 3(3a-1) + \frac{3a(3a-1) - (2a+e)(2a+e-1)}{a+e-1},$$

i.e.

$$\begin{aligned} & (2a+e)(2a+e-1)(a+2e-1) - 3(3a-1)(2a+e-1)e \\ &= 2(2a+e-1)(a-e)(a-e-1) > 0. \end{aligned}$$

Here $a > e + 1$, i.e. $m_0 = 2a + e < d - 1$ since D is indecomposable. So the assertion is proved.

2. Calculation of b -functions

2.1. For a nonconstant polynomial $f \in K[x_1, \dots, x_n]$ with $\text{char } K = 0$, the b -function $b_f(s)$ can be defined to be the minimal polynomial of the action of s on

$$\mathcal{D}_n[s]f^s / \mathcal{D}_n[s]f^{s+1}.$$

This implies that $b_f(s)$ is invariant by extensions of K and its roots are rational numbers since the last assertion holds for $K = \mathbf{C}$ by [Ka1].

Let $i_f : X \hookrightarrow X \times \mathbf{A}_K^1$ denote the graph embedding of f where $X = \mathbf{A}_K^n$. Then via the global section functor, $R_f[s]f^s$ is identified with the direct image by i_f of the $\mathcal{D}_X$ -module $\mathcal{O}_X[\frac{1}{f}]$ in the notation of the introduction. This is compatible with extensions of K . Moreover, the regular holonomic $\mathcal{D}_X$ -module $\text{Gr}_V^\alpha((i_f)_*\mathcal{O}_X[\frac{1}{f}])$ corresponds via the global section functor to $\text{Gr}_V^\alpha(R_f[s]f^s)$, and via the de Rham functor to the λ -eigenspace of Deligne's nearby cycle sheaf $\psi_f \mathbf{C}_X$ ([De]) with $\lambda = e^{-2\pi i \alpha}$ if $K = \mathbf{C}$, see [Ka2], [Ma2].

This implies that the union of the eigenvalues of the Milnor monodromies for $x \in f_{\mathbf{C}}^{-1}(0) \subset \mathbf{C}^n$ is independent of the choice of an embedding $K \hookrightarrow \mathbf{C}$ since the α are rational numbers.

2.2. b -functions of homogeneous polynomials. Assume that $X = \mathbf{C}^n$ and f is a homogeneous polynomial. Let F_f denote the Milnor fiber of f , and $H^{n-1}(F_f, \mathbf{C})_\lambda$ be the λ -eigenspace of the Milnor cohomology by the action of the monodromy T , where $n = \dim X$. Set

$$\lfloor \alpha \rfloor = \max\{k \in \mathbf{Z} \mid k \leq \alpha\}, \quad \mathbf{e}(\alpha) = \exp(2\pi i \alpha) \quad \text{for } \alpha \in \mathbf{Q}.$$

By [Sa1], Th. 2, there is a decreasing filtration P on $H^{n-1}(F_f, \mathbf{C})_\lambda$ such that

$$(2.2.1) \quad b_f(-\alpha) = 0 \quad \text{if} \quad \text{Gr}_P^{\lfloor n-\alpha \rfloor} H^{n-1}(F_f, \mathbf{C})_{\mathbf{e}(-\alpha)} \neq 0,$$

where P coincides with $\tilde{P}$ in loc. cit. since f is homogeneous.

Set $U := \mathbf{P}^{n-1} \setminus Z$ with $Z := f^{-1}(0) \subset \mathbf{P}^{n-1}$. By [Sa1], Prop. 4.9, the filtration P on $H^{n-1}(F_f, \mathbf{C})_\lambda$ is induced by the pole order filtration P on the meromorphic extension $\mathcal{L}^{(k)}$ of a local system $L^{(k)}$ of rank one on U such that

$$(2.2.2) \quad H^j(U, L^{(k)}) = H^j(F_f, \mathbf{C})_\lambda,$$

where $\lambda = \exp(-2\pi i k/d)$ with $d = \deg f$. Here the local system $L^{(k)}$ is defined by the decomposition

$$\pi_* \mathbf{C}_{F_f} = \sum_{k=0}^{d-1} L^{(k)},$$

where π is the canonical projection from the affine Milnor fiber $F_f := f^{-1}(1) \subset \mathbf{C}^n$ onto $U \subset \mathbf{P}^{n-1}$, and the action of the monodromy is the multiplication by $\exp(-2\pi i k/d)$ on $L^{(k)}$ so that (2.2.2) holds, see [CS] or [Di], Prop. 6.4.6. Since $\mathbf{P}^{n-1}$ is simply connected, the local system $L^{(k)}$ is determined by the monodromies around the irreducible components Z_j

of Z . These are given by the multiplication by $\exp(2\pi i m_j k/d)$ where m_j is the multiplicity of the divisor Z along Z_j .

We can identify locally $\mathcal{L}^{(k)}$ with $\mathcal{O}_Y(*Z)h^{-k/d}$ as a $\mathcal{D}_Y$ -module if h defines locally $Z \subset Y := \mathbf{P}^{n-1}$. Then the pole order filtration P on $\mathcal{L}^{(k)}$ is defined by

$$(2.2.3) \quad P_i \mathcal{L}^{(k)} = \mathcal{O}_Y h^{-\frac{k}{d}-i} \quad \text{if } i \geq 0, \text{ and } 0 \text{ otherwise.}$$

Note that the residue of the logarithmic connection on $P_i \mathcal{L}^{(k)}$ at a general point of Z_j is the multiplication by

$$(2.2.4) \quad \left(-\frac{k}{d} - i\right) m_j.$$

The filtration $P^i = P_{-i}$ on $H^{n-1}(U, L^{(k)}) = H^{n-1}(F_f, \mathbf{C})_\lambda$ is induced by P_{n-1-i} on $\mathcal{L}^{(k)}$ using the de Rham complex

$$\mathcal{L}^{(k)} \rightarrow \mathcal{L}^{(k)} \otimes_{\mathcal{O}_Y} \Omega_Y^1 \rightarrow \cdots \rightarrow \mathcal{L}^{(k)} \otimes_{\mathcal{O}_Y} \Omega_Y^{n-1},$$

since the latter has the filtration $P^i = P_{-i}$ defined by

$$P_{-i} \mathcal{L}^{(k)} \rightarrow P_{1-i} \mathcal{L}^{(k)} \otimes_{\mathcal{O}_Y} \Omega_Y^1 \rightarrow \cdots \rightarrow P_{n-1-i} \mathcal{L}^{(k)} \otimes_{\mathcal{O}_Y} \Omega_Y^{n-1}.$$

We have also the Hodge filtration F on $\mathcal{L}^{(k)}$ such that

$$F_i \mathcal{L}^{(k)} \subset P_i \mathcal{L}^{(k)},$$

and the Hodge filtration F on $H^{n-1}(U, L^{(k)}) = H^{n-1}(F_f, \mathbf{C})_\lambda$ is induced by the above formula with P replaced by F .

2.3. Calculation of the cohomology of $L^{(k)}$. From now on, assume $D = f^{-1}(0)$ is a central hyperplane arrangement in $\mathbf{C}^n$. Let D_i ($i = 1, \dots, e$) be the irreducible components of D with multiplicity m_i . Then $Z = \mathbf{P}(D) \subset \mathbf{P}^{n-1}$ and $Z_i = \mathbf{P}(D_i)$. Let D^{nnc} denote the smallest subset of D such that $D \setminus D^{\text{nnc}}$ is a divisor with normal crossings. Set $Z^{\text{nnc}} = \mathbf{P}(D^{\text{nnc}}) \subset \mathbf{P}^{n-1}$. Note that $d = \deg f = \sum_{i=1}^e m_i$.

For $k \in \{0, \dots, d-1\}$ and $I \subset \{1, \dots, e-1\}$ with $|I| = k-1$, define

$$(2.3.1) \quad \begin{aligned} \alpha_i^I &= \begin{cases} -m_i k/d & \text{if } i \notin I \cup \{e\}, \\ 1 - m_i k/d & \text{if } i \in I \cup \{e\}. \end{cases} \\ \alpha_L^I &= \sum_{D_i \supset L} \alpha_i^I. \\ \Sigma^I &= \{p \in Z^{\text{nnc}} \setminus Z_e \mid \alpha_p^I = 0\}, \end{aligned}$$

where L is an edge of D , and we set $\alpha_p^I := \alpha_L^I$ if $\mathbf{P}(L) = \{p\}$. (See Remark 3.6 (iii) below for another way of the definition of the α_i^I .) Here it should be noted that in order to apply the theory in [ESV] (and also in [STV]), we must have a regular singular connection on a *trivial* line bundle, i.e. the following condition should be satisfied:

$$(2.3.2) \quad \sum_{i=1}^e \alpha_i^I = 0.$$

This is satisfied in this case since $d = \sum_{i=1}^e m_i$. Note also that α_e^I is used in an essential way for (2.3.4) below (i.e. the condition of [STV]) although it does not appear in the definition of the connection on the affine space $\mathbf{C}^{n-1}$ which is given below.

For $i \in \{1, \dots, e-1\}$, let $e_i = dg_i/g_i$ with g_i a linear function defining $Z_i \setminus Z_e$ in $\mathbf{P}^{n-1} \setminus Z_e \cong \mathbf{C}^{n-1}$. Set

$$\omega_I := \sum_{i=1}^{e-1} \alpha_i^I e_i.$$

It defines a connection ∇^{ω_I} on $\mathcal{O}_U$ (where $U = \mathbf{P}^{n-1} \setminus Z$) such that

$$\nabla^{\omega_I} u = du + u\omega_I \quad \text{for } u \in \mathcal{O}_U.$$

The corresponding local system is isomorphic to $L^{(k)}$ by comparing their local monodromies as remarked in (2.2). Consider the de Rham cohomology $H_{\text{DR}}^\bullet(U, (\mathcal{O}_U, \nabla^{\omega_I}))$, which is calculated by the complex of rational forms $(\Omega_U^\bullet(U), \nabla^{\omega_I})$ since U is affine. Set

$$\mathcal{A}^p = \sum_{i_1 < \dots < i_p} \mathbf{C} e_{i_1} \wedge \dots \wedge e_{i_p}.$$

Then we have a natural inclusion of complexes

$$(2.3.3) \quad \iota_I^\bullet : (\mathcal{A}^\bullet, \omega_I \wedge) \hookrightarrow (\Omega_U^\bullet(U), \nabla^{\omega_I}),$$

where the source is called the Aomoto complex. Note that we have for $a \in \mathcal{A}^0 = \mathbf{C}$

$$\iota_I^p(a e_{i_1} \wedge \dots \wedge e_{i_p}) = \iota_I^0(a) e_{i_1} \wedge \dots \wedge e_{i_p},$$

and the image of the injection $\iota_I^0 : \mathcal{A}^0 (= \mathbf{C}) \hookrightarrow \Gamma(U, \mathcal{O}_U)$ depends on the choice of I . Indeed, $\text{Im } \iota_I^0$ depends on the trivialization of the line bundle $\mathcal{L}^{(k)}$ which is determined by I , see the proof of Theorem (2.5) and Remark (2.7)(i) below.

By [ESV], [STV], (2.3.3) is a quasi-isomorphism if the following condition is satisfied:

$$(2.3.4) \quad \alpha_L^I \notin \mathbf{Z}_{>0} \text{ for any nonzero dense edges } L \subset D.$$

2.4. Remark. Assume D is reduced (i.e. $m_i = 1$) and $(k, d) = 1$. Then condition (2.3.4) is satisfied for any I with $|I| = k-1$ since $\alpha_L^I \notin \mathbf{Z}$ for any nonzero edge L . Moreover, this assumption implies that $\psi_{f,\lambda} \mathbf{C}_X$, the nearby cycle sheaf with eigenvalue $\lambda := \exp(-2\pi i k/d)$, is supported at the origin. (Indeed, in case the last assertion is not true, there is $d' \in (0, d)$ and $k' \in \mathbf{N}$ such that $k/d = k'/d'$. This follows from the calculation of the Milnor cohomology in (2.2) to $x \in D \setminus \{0\}$. Here the degree d' of the defining equation of D at $x \in D \setminus \{0\}$ becomes strictly smaller. But this contradicts the assumption $(k, d) = 1$.) The above assertion implies further the vanishing of the lower Milnor cohomology $H^j(F_f, \mathbf{C})_\lambda$ for $j < n-1$, since the nearby cycle sheaf $\psi_{f,\lambda} \mathbf{C}_X$ is a perverse sheaf up to the shift of complex by $n-1$. If moreover D is indecomposable, then we get the nonvanishing of the highest Milnor cohomology $H^{n-1}(F_f, \mathbf{C})_\lambda$ by (2.2.2), since the indecomposability is equivalent to the nonvanishing of the Euler characteristic $\chi(U)$, see [STV].

Note that Theorem 4.2(e) in [Sa2] remains valid in the non-reduced case as follows.

2.5. Theorem. *Let $V(I)'$ be the subspace of $\mathcal{A}^{n-1}$ generated by $e_J := e_{j_1} \wedge \cdots \wedge e_{j_{n-1}}$ for any $J = \{j_1, \dots, j_{n-1}\} \subset I$ with $j_1 < j_2 < \cdots < j_{n-1}$. Let $V(I)$ be the image of $V(I)'$ in $H^{n-1}(\mathcal{A}^\bullet, \omega_I \wedge)$, where ω_I and $\alpha^I = (\alpha_i^I)$ are as in (2.3). Assume $V(I) \neq 0$ and (2.3.4) holds. Then $b_f(-\frac{k}{d}) = 0$.*

Proof. By (2.2.1) it is enough to show that the image of e_J by the injection ι_I^{n-1} in (2.3.3) is contained $P_0 \mathcal{L}^{(k)} \otimes_{\mathcal{O}_Y} \Omega_Y^{n-1}$ in the notation of (2.2). Here $P^n H^{n-1}(F_f, \mathbf{C})_\lambda = 0$ since $P_{-1} \mathcal{L}^{(k)} = 0$. By definition the image of $a \in \mathcal{A}^0 = \mathbf{C}$ by ι_I^0 is a global section v_a of a free $\mathcal{O}_Y$ -submodule $\mathcal{L}_I$ of $\mathcal{L}^{(k)}$ such that the residue of the connection at the generic point of Z_i is the multiplication by α_i^I in (2.3.1). Set $Z^{I \cup \{e\}} := \bigcup_{k=1}^n Z_{j_k}$ with $j_n := e$. Then

$$v_a \otimes e_J \in \mathcal{L}_I(Z^{I \cup \{e\}}) \otimes \Omega_Y^{n-1},$$

since $e_J \in \Omega_Y^{n-1}(\log Z^{I \cup \{e\}}) = \Omega_Y^{n-1}(Z^{I \cup \{e\}})$. Thus the assertion is reduced to

$$\mathcal{L}_I(Z^{I \cup \{e\}}) \subset P_0 \mathcal{L}^{(k)},$$

and this is shown by comparing (2.2.4) and (2.3.1). Indeed, the eigenvalue of the residue of the connection on $\mathcal{L}_I(Z^{I \cup \{e\}})$ is shifted by -1 at the generic point of Z_j for $j \in I \cup \{e\}$, but it is not smaller than $-m_j k/d$ even after this shift by (2.3.1). So Theorem (2.5) is proved.

2.6. Proof of Theorem 2 in cases (i) and (iii). In case (i), n/d is a jumping coefficient by Teitler's refinement [Te] of Mustařă's formula [Mu] for multiplier ideals using only dense edges. Hence it is a root of $b_f(s)$ up to a sign by [ELSV].

In case (iii), condition (2.3.4) is satisfied for any I with $|I| = n - 1$ since $k = n$ and $(n, d) = 1$, see Remark (2.4) above. By [FT], Example 4.5, the highest degree cohomology of the Aomoto complex $H^{n-1}(\mathcal{A}^\bullet, \omega_I \wedge)$ has a monomial basis (independently of I) under the genericity condition on D_d . Take a subset

$$I = \{i_1, \dots, i_{n-1}\} \subset \{1, \dots, d-1\},$$

such that the corresponding form $e_I = e_{i_1} \wedge \cdots \wedge e_{i_{n-1}}$ is a member of the obtained monomial basis. Since (2.3.4) is satisfied, the image of e_I in the cohomology of the local system does not vanish. So the assertion follows from Theorem (2.5) (i.e. [Sa2], Th. 4.2(e)).

2.7. Remarks. (i) In the above argument, the image of e_I by ι_I^{n-1} is independent of the choice of I up to a nonzero constant multiple. Indeed, the injection ι_I^0 in (2.3.3) is defined by using the trivial line bundle $\mathcal{L}_I$ in the proof of Theorem (2.5) which is determined by the eigenvalues α_i^I in (2.3.1). If we take another $I' \subset \{1, \dots, d-1\}$ with $|I'| = n - 1$ and $e_{I'} \neq 0$, then, using the trivialization given by $\mathcal{L}_I$, a nonzero constant section of $\mathcal{L}_{I'}$ is identified with the rational function $c g_{I'} / g_I$ where $c \in \mathbf{C}^*$ and $g_I = \prod_{i \in I} g_i$ in the notation of (2.3). This gives the difference between ι_I^j and $\iota_{I'}^j$ for any j . So the independence follows since $g_I e_I = c' g_{I'} e_{I'}$ with $c' \in \mathbf{C}^*$.

(ii) We can also identify the image of e_I by ι_I^{n-1} with an element of the Gauss-Manin system of f . The problem is then closely related to the torsion of the Brieskorn lattice.

3. The rank 3 case

In this section we assume $n = 3$ and give two proofs of the case (ii) in Theorem 2. Note that the case $n \leq 2$ is well-known. Indeed, it follows for instance from [Mu], [ELSV].

3.1. Conditions. From now on we assume

$$n = k = 3.$$

We will write $p \subset i$ if $\{p\} \subset Z_i$, and set $\alpha_p^I = \alpha_L^I$ if $\mathbf{P}(L) = \{p\}$.

In the notation of (2.3.1) we will study the following three conditions:

- (a) $\alpha_p^I \notin \mathbf{Z}_{>0}$ for any $p \in Z^{\text{nnc}} = \mathbf{P}(D^{\text{nnc}})$.
- (b) $\exists p_0 \in (\bigcup_{i \in I} Z_i)^{\text{sing}} \setminus Z_e$.
- (c) $Z \setminus (Z_e \cup \Sigma^I \cup \{p_0\})$ is connected.

3.2. Remarks. (i) In the case $n = 3$, condition (a) coincides with condition (2.3.4) which implies that the inclusion (2.3.3) is a quasi-isomorphism. Note that we have always the inequality of the dimensions, see [LY], Prop. 4.2.

(ii) For $i, j, k \supset p$, there is a well-known relation

$$(3.2.1) \quad e_i \wedge e_j = e_i \wedge e_k - e_j \wedge e_k,$$

which is easily checked by setting $g_i = x$, $g_j = y$ and $g_k = x + y$. This also follows from the relations of the Orlik-Solomon algebra which are given by $\partial(e_i \wedge e_j \wedge e_k)$ for $i, j, k \supset p$, see e.g. [OT], p. 60. As in [BDS], Lemma 1.4, this implies for $\eta = \sum_{i=1}^{e-1} \beta_i e_i$ and $p \in Z^{\text{nnc}} \setminus Z_e$

$$(3.2.2) \quad \text{If } \pi_p(\omega_I \wedge \eta) = 0, \text{ then } \alpha_p^I \beta_i = \beta_p \alpha_i^I \text{ for any } i \supset p.$$

Here $\beta_p = \sum_{i \supset p} \beta_i$, and $\pi_p(\omega_I \wedge \eta)$ is the p -component in the direct sum decomposition in [BDS], 1.3.2

$$H^2(U, \mathbf{Q}) = \bigoplus_p L_p,$$

where p runs over $(Z_{\text{red}})_{\text{sing}} \setminus Z_e$, and L_p is a vector space of rank $m'_p - 1$ with m'_p the multiplicity of Z_{red} at p . More precisely L_p has a basis consisting of $e_i \wedge e_k$ with $i \supset p$ and $i \neq k$ where k is any fixed member such that $k \supset p$. This also follows from the definition of the Orlik-Solomon algebra mentioned after (3.2.1), see e.g. [OT], p. 60.

We also get

$$(3.2.3) \quad \text{If } p \in (Z_i \cap Z_j) \setminus (Z^{\text{nnc}} \cup Z_e), \text{ then } \alpha_i^I \beta_j = \alpha_j^I \beta_i.$$

In case $\alpha_i^I \neq 0$ (i.e. $m_i \neq d/3$) for any $i \in I$, we have by (3.2.2–3)

$$(3.2.4) \quad \text{If } \pi_p(\omega_I \wedge \eta) = 0 \text{ and } p \notin \Sigma^I, \text{ then } \beta_i/\alpha_i^I \text{ is independent of } i \supset p.$$

(iii) Lemma 1.4 in [BDS] or above (3.2.2) is essentially known to the specialists, see [LY], Lemma 3.1 (and also [Fa], [Li2], [Yu]). Here the situation is localized at p , i.e. the lines not passing through p are neglected, by using the fact that the relations of the Orlik-Solomon algebra are of the form $\partial(e_J)$ for certain J and are compatible with the decomposition by p .

3.3. Proposition. *With the notation and the assumption of (2.3), assume $n = k = 3$ and there is $I \subset \{1, \dots, e-1\}$ such that $|I| = 2$ and conditions (a), (b) and (c) in (3.1) are satisfied. Then $b_f(-3/d) = 0$ where f is a defining polynomial of D .*

Proof. Let p_0 be as in condition (b) in (3.1), and assume the following condition is satisfied:

$$\pi_p(\omega_I \wedge \eta) = 0 \text{ for any } p \neq p_0.$$

Then η is a multiple of ω_I , i.e. β_i/α_i^I is independent of i , see Remark (3.2)(ii). So we can apply Theorem (2.5) (i.e. [Sa2], Th. 4.2(e)), and conclude that $b_f(-3/d) = 0$. This finishes the proof of Proposition (3.3).

3.4. One proof of Theorem 2(ii). We may assume that $\{0\}$ is not a good dense edge, since we can apply the case (i) otherwise. By Proposition (3.3), it is sufficient to show the following:

Assertion. There is an irreducible component Z_e of Z together with a subset $I \subset \{1, \dots, e-1\}$ such that $|I| = 2$ and conditions (a), (b) and (c) in (3.1) are satisfied changing the order of $\{1, \dots, e\}$ if necessary.

Note first that α_p^I can be an integer only in the case $d/3 \in \mathbf{Z}$. (Indeed, we have $m_p := \sum_{i \supset p} m_i < d$, and hence $\alpha_p^I \equiv 3m_p/d \not\equiv 0 \pmod{\mathbf{Z}}$ unless $d/3 \in \mathbf{Z}$.) Then the above assertion is shown in the case $d/3 \notin \mathbf{Z}$ as follows.

Since $\alpha_p^I \notin \mathbf{Z}$ for any $p \in Z^{\text{nnc}}$, condition (a) is trivially satisfied and $\Sigma^I = \emptyset$ for any choice of I . Assuming D central and indecomposable, there is $p_0 \in Z^{\text{sing}}$ together with Z_e and I satisfying condition (b). As for condition (c), it is not satisfied only in the case there is Z_i passing through p_0 and such that $Z_i \cap Z_{i'} \subset (Z_i \cap Z_e) \cup \{p_0\}$ for any $i' \notin \{i, e\}$. (Otherwise, for any Z_i passing through p_0 , there is $Z_{i'}$ such that $Z_i \cap Z_{i'} \not\subset Z_e \cup \{p_0\}$.) In this case every Z_i passes through either p_0 or $Z_i \cap Z_e$. This implies that $|Z^{\text{nnc}}| = 2$ since D is indecomposable. Then, replacing Z_e with Z_i containing Z^{nnc} , we may take p_0 to be any point of $Z^{\text{sing}} \setminus Z^{\text{nnc}}$ and I is chosen so that $\{p_0\} = \bigcap_{i \in I} Z_i$. Thus the assertion is proved in this case.

We may now assume

$$a := d/3 \in \mathbf{Z}.$$

Since $\{0\}$ is not a good dense edge, there is $p_1 \in Z^{\text{nnc}}$ with multiplicity $> 2a$. On the other hand, we may assume that there is $p_2 \in Z^{\text{nnc}}$ with $\alpha_{p_2}^I \in \mathbf{Z}$, i.e. its multiplicity

is divisible by a , since otherwise the above conditions are easily satisfied. Thus we may assume that there are $p_1, p_2 \in Z^{\text{unc}}$ with multiplicity $2a + 1$ and a respectively and hence $Z^{\text{unc}} = \{p_1, p_2\}$, since $d = 3a$. So the assertion is proved by the same argument as above.

3.5. Another proof of Theorem 2(ii). It is also possible to prove Theorem 2(ii) by taking p_0 to be the point with multiplicity $m_{p_0} > \frac{2}{3}d$, which exists since we may assume that $\{0\}$ is not a good dense edge as in (3.4). In this case there is a line Z_{d-1} which is different from the line at infinity Z_d and does not contain p_0 since D is indecomposable. Moreover there are at least two lines Z_1, Z_2 passing through p_0 such that their intersections with Z_{d-1} are ordinary double points of Z and furthermore their intersections with Z_d do not have multiplicity a so that conditions (a) and (b) in (3.1) are satisfied by setting $I = \{1, 2\}$. Indeed, we have $m_{p_0} > \frac{2}{3}d$, $d - m_{p_0} \geq 2$, and hence $d > 6$, and moreover the number of lines Z_i such that $i \supset p_0$ and $Z_i \cap Z_{d-1}$ is an ordinary double point of Z is at least

$$m_{p_0} - 1 - (d - 2 - m_{p_0}) > 1,$$

since $|\bigcup_{i \supset p_0} Z_i \cap Z_{d-1}| \geq m_{p_0} - 1$. So the condition on the intersection with Z_{d-1} is satisfied. For the intersection with Z_d we can exclude the case where a point of Z has multiplicity a since this case has a very special structure as explained at the end of (3.4) (e.g. the singular points of Z other than this point and p_0 are ordinary double points) so that we can easily choose Z_1, Z_2 satisfying the above conditions in this case.

We can then prove Theorem 2(ii) without using Proposition (3.3) but using (3.2.1). Indeed, by Theorem (2.5) (i.e. [Sa2], Th. 4.2(e)), it is enough to show

$$(3.5.1) \quad \text{If } (\sum_i \alpha_i^I e_i) \wedge (\sum_j \beta_j e_j) = c e_1 \wedge e_2 \text{ for some } c \in \mathbf{Q}, \text{ then } c = 0.$$

Under the assumption of (3.5.1) we get by using (3.2.1)

$$(3.5.2) \quad \alpha_{p_0}^I \beta_i = \beta_{p_0} \alpha_i^I \text{ if } i \supset p_0 \text{ and } i > 2.$$

Here we have $\alpha_{p_0}^I \neq 0$ since $m_{p_0} > \frac{2}{3}d$. So we may assume

$$(3.5.3) \quad \beta_i = 0 \text{ if } i \supset p_0 \text{ and } i > 2,$$

by replacing β_i with $\beta_i - c' \alpha_i^I$ for any i where $c' := \beta_{p_0} / \alpha_{p_0}^I$. (Note that this change of β_i does not affect the hypothesis of (3.5.1).) Since $m_{p_0} > 4$, (3.5.2) and (3.5.3) imply

$$\beta_1 + \beta_2 = \beta_{p_0} = 0.$$

On the other hand, by (3.2.3) applied to the intersections of Z_1, Z_2 with Z_{d-1} , we get

$$\beta_1 / \alpha_1^I = \beta_{d-1} / \alpha_{d-1}^I = \beta_2 / \alpha_2^I,$$

where $\alpha_1^I = \alpha_2^I \neq 0$ and $\alpha_{d-1}^I \neq 0$ since Z is reduced. So $\beta_1 = \beta_2 = 0$, and (3.5.1) follows.

3.6. Remarks. (i) It does not seem easy to generalize the above arguments to the non-reduced case. If p_0 is taken to be the point with the highest multiplicity, there is an example as follows: Assume $a > 6$, and let

$$f = (xy(x-y))^{a-2}(x+y-z)(x+y-2z)(x+2y-2z)(2x+y-2z)z^2.$$

Here $d = 3a$, and there does not exist I such that the argument in (3.5) can be applied if we set $p_0 = (0, 0, 1)$. Indeed, let Z_i ($i = 1, \dots, 8$) denote the lines defined by the linear factors of f respecting the order of the factors, where $e = 8$. Here Z_e must be the line defined by $z = 0$ since conditions (a) and (b) in (3.1) cannot be satisfied otherwise. Then the singular points of $Z \setminus (\{p_0\} \cup Z_e)$ contained in Z_1 or Z_2 have all multiplicity a , and moreover Z_{red} has multiplicity 3 at these points. So the argument in (3.5) cannot be applied.

(ii) For a more complicated example, we might consider the following: Let E be an elliptic curve in the dual projective space $\mathbf{P}^2$, and G be the subgroup of torsion points of order three. This defines a projective hyperplane arrangement in $\mathbf{P}^2$ with $e = |G| = 9$, see e.g. [Li]. Let G_0 be a subgroup of G with order 3. Assume $a > 6$. To the lines corresponding to the elements of G_0 we give the multiplicity $a - 2$, while the other lines have multiplicity 1. Then $d = 3a$, and $I \cup \{e\}$ should correspond to $G_0 + p \subset G$ for some $p \in G$ in order to satisfy condition (a) in (3.1). (Indeed, if there are $g_1, g_2 \in I \cup \{e\}$ such that their images in G/G_0 are different, then there is $g_3 \in G$ such that the images of g_1, g_2, g_3 in G/G_0 are all different and moreover $g_1 + g_2 + g_3 = 0$. The last condition is equivalent to the condition that the three lines corresponding to g_1, g_2, g_3 intersect at one point. Then condition (a) is not satisfied at this point.) So p_0 is contained in Z_e , and hence condition (b) cannot be satisfied. Thus we cannot prove a generalization of Theorem 2 in this case by using Theorem (2.5) (i.e. the generalization of [Sa2], Th. 4.2(e) to the nonreduced case).

(iii) In order to apply the theory in [ESV] and [STV], we have to choose the residues α_i of the connection satisfying the two conditions (2.3.2) and (2.3.4). In our case we have $\alpha_i = n_i - m_i k/d$ with $n_i \in \mathbf{Z}$ by the monodromy condition, and $\sum_i n_i = k$ since $\sum_i m_i = d$. Then, to satisfy (2.3.2), an easy way is to choose a subset J of $\{1, \dots, e\}$ with $|J| = k$ and set $n_i = 1$ for $i \in J$ and $n_i = 0$ otherwise. Here there are two possibilities depending on whether $e \in J$ or $e \notin J$. Since e corresponds to the divisor at infinity, this makes some difference in the calculation of the Aomoto complex which is defined on the complement affine space $\mathbf{C}^{n-1}$. In (2.3.1) we considered the former case where $I = J \setminus \{e\}$. However, it is also possible to consider the latter case where $I = J$ so that $|I| = k$ instead of $|I| = k - 1$, and

$$\alpha_i^I = \begin{cases} -m_i k/d & \text{if } i \notin I, \\ 1 - m_i k/d & \text{if } i \in I. \end{cases}$$

In the latter case, however, it is usually more difficult to satisfy the three conditions in (3.1).

(iv) If $n = 3$, $d \leq 7$ and $\text{mult}_p Z = 3$ for any $p \in Z^{\text{nnc}}$ in the notation of (2.3), the b -function of a reduced hyperplane arrangement is calculated in [Sa2].

Appendix

by Willem Veys

University of Leuven, Department of Mathematics
Celestijnenlaan 200 B, B-3001 Leuven (Heverlee), Belgium

This appendix describes some examples solving Question (Q) in (1.4) negatively. I thank the authors of this paper for writing some details.

A.1. Example. We first explain an example of a nonreduced hyperplane arrangement with $n = 3$, $d = 9$. Let

$$f = xy(x - y)z^2(x - z)^4.$$

This gives a negative answer to Question (Q) in (1.4). Indeed, we have $\chi(U) = 1$ by using the affine space defined by $x \neq 0$. We have $\chi(Z_i^\circ) = -1$ except for the line defined by $x = 0$, and the Euler characteristic is 0 for the latter. So we get

$$\begin{aligned} Z_{f,0}^{\text{top}}(s) = & \frac{1}{9s+3} \left(1 - \frac{2}{s+1} - \frac{1}{2s+1} - \frac{1}{4s+1} + \left(-1 + \frac{3}{s+1} \right) \frac{1}{3s+2} \right. \\ & \left. + \left(-1 + \frac{1}{s+1} + \frac{1}{2s+1} + \frac{1}{4s+1} \right) \frac{1}{7s+2} + \frac{2}{s+1} \left(\frac{1}{2s+1} + \frac{1}{4s+1} \right) \right). \end{aligned}$$

Set $\Phi(s) = (9s+3)Z_{f,0}^{\text{top}}(s)$. Since $\frac{1}{2s+1} + \frac{1}{4s+1}$ vanishes by substituting $s = -1/3$, we get

$$\Phi(-1/3) = 1 - 3 + \left(-1 + \frac{9}{2} \right) - 3 \left(-1 + \frac{3}{2} \right) = 0.$$

So the pole at $-1/3$ vanishes. (It vanishes also for the motivic or Hodge zeta function.)

Note that the above example does not give a counterexample to Conjecture (C). This is shown by using Theorem (2.5) and Remark (3.2)(ii) below. Here $p = (0 : 1 : 0) \in \mathbf{P}^2$, the line at infinity is $\{y = 0\}$, and I corresponds to the two lines with multiplicities 2 and 4. This assertion is also shown by a calculation using the computer program Asir.

A.2. Example. There is an example of a *reduced* hyperplane arrangement with $n = 5$ and $d = 10$, giving a negative answer to Question (Q) in (1.4), and which is defined by a polynomial f as below:

$$f = (x - y)(x - 2y)(x - 3y)(x - 4y)(x - 5y)(x + y + z)zuv(u + v + z).$$

In fact, let Z_1, Z_2, Z_3 be closed subvarieties of $Y := \mathbf{P}^4$ defined by

$$Z_1 = \{x = y = z = 0\}, \quad Z_2 = \{u = v = z = 0\}, \quad Z_3 = \{x = y = 0\}.$$

Let $\rho : Y' \rightarrow Y$ be the composition of the blow-up of Y along Z_1, Z_2 and the blow-up along the proper transform of Z_3 . This gives an embedded resolution of (Y, Z) where $Z := \{f = 0\} \subset Y$. We have a partition $\{S_i\}_{i=0,\dots,3}$ of $Y = \mathbf{P}^4$ defined by

$$S_0 = \{z \neq 0\}, \quad S_i = Z_i \ (i = 1, 2), \quad S_3 = \{z = 0\} \setminus (Z_1 \cup Z_2).$$

Consider the pullback of the partition

$$S'_i := \rho^{-1}(S_i) \quad (i = 0, \dots, 3).$$

Let x', y', u', v' be affine coordinates of S_0 defined respectively by $\frac{x}{z}, \frac{y}{z}, \frac{u}{z}, \frac{v}{z}$. Then

$$S'_0 = \tilde{\mathbf{C}}_{x',y'}^2 \times \mathbf{C}_{u',v'}^2, \quad S'_1 = \tilde{\mathbf{P}}_{x,y,z}^2 \times \mathbf{P}_{u,v}^1, \quad S'_2 = \mathbf{P}_{u,v,z}^2 \times \mathbf{P}_{x,y}^1,$$

where $\tilde{\mathbf{C}}_{x',y'}^2$ and $\tilde{\mathbf{P}}_{x,y,z}^2$ are respectively the blow-up of $\mathbf{C}_{x',y'}^2$ and $\mathbf{P}_{x,y,z}^2$ along $(0,0)$ and $(0:0:1)$. Here the lower indices x,y etc. indicate the coordinates. Note that each S_i is a union of strata of the stratification associated to the divisor with normal crossings $\rho^{-1}(Z)$. So we get

$$Z_{f,0}^{\text{top}}(s) = \sum_{i=0}^3 \frac{\Psi_i(s)}{10s+5},$$

where $\Psi_i(s)/(10s+5)$ is the sum of the factors of $Z_{f,0}^{\text{top}}(s)$ associated to the strata contained in S'_i . Since the stratification is compatible with the above product structure, we get

$$\begin{aligned} \Psi_0(s) &= \left(4 - \frac{9}{s+1} + \frac{5}{(s+1)^2} + \left(-3 + \frac{5}{s+1}\right) \frac{1}{5s+2}\right) \cdot \left(1 - \frac{3}{s+1} + \frac{3}{(s+1)^2}\right), \\ \Psi_1(s) &= \frac{1}{7s+3} \left(4 - \frac{13}{s+1} + \frac{11}{(s+1)^2} + \left(-3 + \frac{5}{s+1}\right) \frac{1}{5s+2}\right) \cdot \left(-1 + \frac{3}{s+1}\right), \\ \Psi_2(s) &= \frac{1}{4s+3} \left(1 - \frac{4}{s+1} + \frac{6}{(s+1)^2}\right) \cdot \left(-4 + \frac{6}{s+1}\right), \\ \Psi_3(s) &= 0. \end{aligned}$$

Indeed, let Z'_0 be the divisor on $\mathbf{P}_{x,y,z}^2$ defined by the product of linear factors of f which are linear combinations of x, y, z , and similarly for Z''_0 with x, y replaced by u, v . Then

$$\chi(\mathbf{P}^2 \setminus Z'_0) = 4, \quad \chi(\mathbf{P}^2 \setminus Z''_0) = 1, \quad \chi(Z'_0 \setminus \text{Sing } Z'_0) = -13, \quad \chi(Z''_0 \setminus \text{Sing } Z''_0) = -4,$$

and the number of ordinary double points of Z'_0 and Z''_0 are respectively 11 and 6. The calculation for $\mathbf{P}_{u,v}^1$ and $\mathbf{P}_{x,y}^1$ is similar, and we get the formulas for $\Psi_1(s)$ and $\Psi_2(s)$ since the definition of $\Psi_1(s), \Psi_2(s)$ is compatible with the above product structure using the formula: $\chi(X_1 \times X_2) = \chi(X_1) \cdot \chi(X_2)$ for topological spaces X_1, X_2 . As for the first terms, note that the codimensions of the centers Z_1, Z_2 are 3, and the multiplicities of f at the generic points of Z_1 and Z_2 are respectively 7 and 4. The term $\left(-3 + \frac{5}{s+1}\right) \frac{1}{5s+2}$ comes from the exceptional divisor of the blow-up along the proper transform of Z_3 , where the multiplicity of f at the generic point of Z_3 is 5 and Z_3 has codimension 2.

The argument is similar for $\Psi_0(s)$. Here the Euler number of the smooth part and the number of ordinary double points change since the varieties are restricted to (the blow-up of) the affine space $\mathbf{C}^2$. The vanishing of $\Psi_3(s)$ follows from the $\mathbf{C}^*$ -action on $S'_3 = S_3$ compatible with the stratification, which is defined by $\lambda(x : y : u : v) = (\lambda x : \lambda y : u : v)$ for $\lambda \in \mathbf{C}^*$.

Substituting $s = -\frac{1}{2}$ to the above formulas, we get

$$\Psi_0\left(-\frac{1}{2}\right) = -8 \cdot 7, \quad \Psi_1\left(-\frac{1}{2}\right) = -2 \cdot 8 \cdot 5, \quad \Psi_2\left(-\frac{1}{2}\right) = 17 \cdot 8,$$

and hence the pole of $Z_{f,0}^{top}(s)$ at $s = -\frac{1}{2}$ vanishes. For the moment it is not clear whether $-\frac{1}{2}$ is a root of $b_f(s)$.

References

- [ACLM1] Artal Bartolo, E., Cassou-Noguès, P., Luengo, I. and Melle Hernández, A., Monodromy conjecture for some surface singularities, *Ann. Sci. Ecole Norm. Sup. (4)* 35 (2002), 605–640.
- [ACLM2] Artal Bartolo, E., Cassou-Noguès, P., Luengo, I. and Melle Hernández, A., Quasi-ordinary power series and their zeta functions, *Mem. Amer. Math. Soc.* 178 (2005), no. 841.
- [Be] Bernstein, I.N., Analytic continuation of generalized functions with respect to a parameter, *Funk. Anal.* 6 (1972), 26–40.
- [BDS] Budur, N., Dimca, A. and Saito, M., First Milnor cohomology of hyperplane arrangements, in “Topology of Algebraic Varieties and Singularities”, *Contemporary Mathematics* 538 (2011), pp. 279–292.
- [BMS] Budur, N., Mustașă, M. and Saito, M., Bernstein-Sato polynomials of arbitrary varieties, *Compos. Math.* 142 (2006), 779–797.
- [BMT] Budur, N., Mustașă, M. and Teitler, Z., The monodromy conjecture for hyperplane arrangements, (arXiv:0906.1991) to appear in *Geom. Dedicata*.
- [BS] Budur, N. and Saito, M., Jumping coefficients and spectrum of a hyperplane arrangement, *Math. Ann.* 347 (2010), 545–579.
- [CS] Cohen, D.C. and Suciu, A., On Milnor fibrations of arrangements, *J. London Math. Soc.* 51 (1995), 105–119.
- [DP] De Concini, C. and Procesi, C., Wonderful models of subspace arrangements, *Selecta Math. (N.S.)* 1 (1995), 459–494.
- [De] Deligne, P., Le formalisme des cycles évanescents, in *SGA7 XIII and XIV, Lect. Notes in Math.* 340, Springer, Berlin, 1973, pp. 82–115 and 116–164.
- [Den] Denef, J., Report on Igusa’s local zeta function, *Séminaire Bourbaki* no. 741, *Astérisque* 201/202/203 (1991), 359–386.
- [DL] Denef, J. and Loeser, F., Caractéristiques d’Euler-Poincaré, fonctions zéta locales et modifications analytiques, *J. Amer. Math. Soc.* 5 (1992), 705–720.
- [Di] Dimca, A., *Sheaves in Topology*, Universitext, Springer, Berlin, 2004.
- [ELSV] Ein, L., Lazarsfeld, R., Smith, K.E. and Varolin, D., Jumping coefficients of multiplier ideals, *Duke Math. J.* 123 (2004), 469–506.
- [ESV] Esnault, H., Schechtman V. and Viehweg, E., Cohomology of local systems on the complement of hyperplanes, *Inv. Math.* 109 (1992), 557–561.
- [Fa] Falk, M., Arrangements and cohomology, *Ann. Combin.* 1 (1997), 135–157.
- [FT] Falk, M. and Terao, H., β -bases for cohomology of local systems on hyperplane complements, *Trans. Amer. Math. Soc.* 349 (1997), 189–202.
- [HMY] Howald, J., Mustașă, M. and Yuen, C., On Igusa zeta functions of monomial ideals, *Proc. Amer. Math. Soc.* 135 (2007), 3425–3433.

- [Ig1] Igusa, J.-i., Complex powers and asymptotic expansions, I and II, *J. reine angew. Math.* 268/269 (1974), 110–130 and 278/279 (1975), 307–321.
- [Ig2] Igusa, J.-i., b -functions and p -adic integrals, in *Algebraic analysis*, Vol. I, Academic Press, Boston, MA, 1988, pp. 231–241.
- [Ig3] Igusa, J.-i., Local zeta functions of certain prehomogeneous vector spaces, *Amer. J. Math.* 114 (1992), 251–296.
- [Ig4] Igusa, J.-i., *An introduction to the theory of local zeta functions*, AMS/IP Studies in Advanced Mathematics, 14, American Mathematical Society, Providence, RI; International Press, Cambridge, MA, 2000.
- [Ka1] Kashiwara, M., B -functions and holonomic systems, *Inv. Math.* 38 (1976/77), 33–53.
- [Ka2] Kashiwara, M., Vanishing cycle sheaves and holonomic systems of differential equations, *Lect. Notes in Math.* 1016, Springer, Berlin, 1983, pp. 134–142.
- [KSZ] Kimura, T., Sato, F. and Zhu, X.-W., On the poles of p -adic complex powers and the b -functions of prehomogeneous vector spaces, *Amer. J. Math.* 112 (1990), 423–437.
- [LVa] Lemahieu, A. and Van Proeyen, L., Monodromy conjecture for nondegenerate surface singularities, to appear in *Trans. A.M.S.*
- [LV1] Lemahieu, A. and Veys, W., On monodromy for a class of surfaces, *C. R. Math. Acad. Sci. Paris* 345 (2007), 633–638.
- [LV2] Lemahieu, A. and Veys, W., Zeta functions and monodromy for surfaces that are general for a toric idealistic cluster, *Int. Math. Res. Not.* 2009, 11–62.
- [Li] Libgober, A., Characteristic varieties of algebraic curves, in *Applications of algebraic geometry to coding theory, physics and computation* (Eilat, 2001), NATO Sci. Ser. II Math. Phys. Chem., 36, Kluwer Acad. Publ., Dordrecht, 2001, pp. 215–254.
- [LY] Libgober, A. and Yuzvinsky, S., Cohomology of the Orlik-Solomon algebras and local systems, *Compos. Math.* 121 (2000), 337–361.
- [Lo1] Loeser, F., Fonctions d’Igusa p -adiques et polynômes de Bernstein, *Amer. J. Math.* 110 (1988), 1–21.
- [Lo2] Loeser, F., Fonctions d’Igusa p -adiques, polynômes de Bernstein, et polyèdres de Newton, *J. reine angew. Math.* 412 (1990), 75–96.
- [Ma1] Malgrange, B., Le polynôme de Bernstein d’une singularité isolée, in *Lect. Notes in Math.* 459, Springer, Berlin, 1975, pp. 98–119.
- [Ma2] Malgrange, B., Polynôme de Bernstein-Sato et cohomologie évanescence, *Analysis and topology on singular spaces*, II, III (Luminy, 1981), *Astérisque* 101–102 (1983), 243–267.
- [Mu] Mustață, M., Multiplier ideals of hyperplane arrangements, *Trans. Amer. Math. Soc.* 358 (2006), 5015–5023.
- [OT] Orlik, P. and Terao, H., *Arrangements of hyperplanes*, Springer, Berlin, 1992.
- [Ro] Rodrigues, B., On the monodromy conjecture for curves on normal surfaces, *Math. Proc. Cambridge Philos. Soc.* 136 (2004), 313–324.
- [Sa1] Saito, M., Multiplier ideals, b -function, and spectrum of a hypersurface singularity, *Compos. Math.* 143 (2007) 1050–1068.
- [Sa2] Saito, M., Bernstein-Sato polynomials of hyperplane arrangements (math.AG/0602527).
- [STV] Schechtman, V., Terao, H. and Varchenko, A., Local systems over complements of hyperplanes and the Kac-Kazhdan conditions for singular vectors, *J. Pure Appl. Algebra* 100 (1995), 93–102.

- [Te] Teitler, Z., A note on Mustață's computation of multiplier ideals of hyperplane arrangements, Proc. Amer. Math. Soc. 136 (2008), 1575–1579.
- [VP] Van Proeyen, L., Local zeta functions for ideals and the monodromy conjecture, thesis 2008 Katholieke Universiteit Leuven.
- [VV] Van Proeyen, L. and Veys, W., The monodromy conjecture for zeta functions associated to ideals in dimension two, Annales Inst. Fourier, 60 (2010), 1347–1362.
- [Ve1] Veys, W., On the poles of Igusa's local zeta function for curves, J. London Math. Soc. (2) 41 (1990), 27–32.
- [Ve2] Veys, W., Poles of Igusa's local zeta function and monodromy, Bull. Soc. Math. France 121 (1993), 545–598.
- [Ve3] Veys, W., Determination of the poles of the topological zeta function for curves, Manuscripta Math. 87 (1995), 435–448.
- [Ve4] Veys, W., Vanishing of principal value integrals on surfaces, J. Reine Angew. Math. 598 (2006), 139–158.
- [Yu] Yuzvinsky, S., Cohomology of the Brieskorn-Orlik-Solomon algebras, Comm. Algebra 23 (1995), 5339–5354.

Department of Mathematics, University of Notre Dame, IN 46556, USA

RIMS Kyoto University, Kyoto 606-8502 Japan

Department of Mathematics, University of Oregon, Eugene, Oregon 94703, USA

Mar. 22, 2011, v.3g